ON SOME ARITHMETIC PROPERTIES OF FINITE GROUPS

A.N. Skiba

Francisk Skorina Gomel State University
104 Sovetskaya str., 246019, Gomel, Belarus
alexander.skiba49@gmail.com

We fix some partition $\sigma = \{\sigma_i | i \in I\}$ of the set of all primes $\mathbb{P}$ (that is, $\mathbb{P} = \cup_{i\in I}\sigma_i$ and $\sigma_i \cap \sigma_j = \emptyset$ for all $i \neq j$). A group G is called **σ-primary** if G is a σ_i-group for some $i = i(G)$.

We say that a finite group G is: **σ-soluble** if every chief factor of G is σ-primary; **σ-nilpotent** if $(H/K) \rtimes (G/C_G(H/K))$ is σ-primary for every chief factor H/K of G.

Based on these concepts, we develop and unify [1–5] some aspects of the theories of soluble and quasinilpotent groups, of the subgroup lattices theory and of the theory of subnormal subgroups.

References

1. Skiba A.N. *On σ-subnormal and σ-permutable subgroups of finite groups* // J. Algebra. 2015. V. 436. P. 1–16.

2. Guo W., Skiba *A.N. Finite groups with generalized Ore supplement conditions for primary subgroup* // J. Algebra. 2015. V. 432. P. 205–227.

3. Skiba A.N. *A generalization of a Hall theorem* // J. Algebra and its Application (accepted).

4. Skiba A.N. *On the generalized σ-Fitting subgroup of finite groups* // Proc. Amer. Math. Soc. (submitted).

5. Guo W., Skiba A.N. *Finite groups with permutable complete Wielandt sets of subgroups* // J. Group Theory. 2015. V. 18. No. 2. P. 191–200.